

Christopher T. Donaldson, MD

813.684.BONE (2663)

www.omgtb.com

Post-op PATIENT Instructions: PROXIMAL HUMERUS ORIF

<u>Dressing</u>: A dressing has been applied to your shoulder to absorb any fluid/blood. A small amount of blood on the dressing is expected. Your nurse will teach you how to change your dressing. Change the dressing each day, <u>leaving the steri-strips on</u>. Unless directed by your surgeon, <u>no</u> salves, balms, or ointments (even antibiotic ointments) to the incisions. Soreness and bruising is expected for several days afterward. Showering is permitted 72 hours after surgery with the incisions covered. After showering, apply a new dry dressing. Do NOT soak the incisions in a bathtub. No swimming/hot-tubs/lakes or saltwater until cleared by your surgeon.

<u>Ice</u>: Ice is a powerful anti-inflammatory. Icepacks/wraps will help to reduce swelling and pain. Use liberally (20-30 min./session), but protect the skin from direct contact (and frostbite).

<u>Activity</u>: **NO ACTIVE ATTEMPTING TO LIFT THE ARM ** Wear the immobilizer/sling as directed below. The forearm strap may be removed for hand, wrist, and elbow range of motion which decreases extremity swelling. Sleeping in a recliner with pillows behind the elbow to keep the arm across the belly may provide additional comfort. No driving until permission is given by your surgeon.

<u>Pain</u>: A nerve block has been performed for immediate post-op pain control by the anesthesiologist. It typically "wears off" at about 8-12 hrs following surgery. A narcotic (taken every 3-4hrs as needed for breakthrough pain) is given to control your pain. <u>Begin taking these pain medications when you BEGIN experiencing pain!</u> These meds can take 30-45 minutes to start "working". You do not want to play "catch-up" by letting your pain get out of control. Nausea, drowsiness, and constipation are common side effects of narcotics. Adequate fluid intake and a stool softener obtained over the counter from your local pharmacy is recommended to minimize constipation. Call the office if you are unable to tolerate your medication.

<u>Precautions</u>: Call the office (813) 684-2663 if you develop: temperatures >101°F, shortness of breath, chest pain, uncontrolled pain, marked redness/hives, persistent drainage, new onset numbness, significant incisional/calf swelling, or any other concerns.

Post-operative Appointment:

o <u>Call (813) 684-2663 today to make a post-operative appointment to see your surgeon 10-14 days following your surgery.</u>

Special Instructions: (Additionally, follow any indicated instructions below.)

Physical therapy: prescription will be given at the 1st post op visit

☑ Sling / Shoulder Brace: ☑ for 6 weeks; (NO DRIVING WHILE IN SLING!)

☑ <u>Ice (Cryotherapy) Unit:</u> ☑ protect/check skin regularly

**A blood thinner to prevent blood clots should be taken for 6 weeks following surgery **