



PATIENT NAME:

DOB:

PHARMACY:

POSTOPERATIVE INSTRUCTIONS

PROCEDURE: QUADRICEPS OR PATELLAR TENDON REPAIR

NEXT APPOINTMENT:

LOCATION:

PAIN MEDICATION

You will be given a prescription for pain medication after surgery. This should be filled and ready for use when you return home from surgery. You **SHOULD NOT** drive, operate heavy machinery, or participate in activities that require concentration while taking narcotic pain medications such as Percocet, Oxycodone, Norco, Vicodin, or Tylenol with Codeine. You **SHOULD** take an over-the-counter stool softener while taking narcotic pain medications to prevent constipation.

DRESSING

Leave the dressing intact until you return for your postoperative appointment. Keep dressing clean and dry. **DO NOT** get the incision(s) or dressings wet. You will not be able to soak the incision(s) for a **MINIMUM OF 4 WEEKS** after surgery.

BRACE

You were placed in a knee brace after surgery. This brace should be worn **AT ALL TIMES**, including when sleeping. It **MUST BE** locked in extension **AT ALL TIMES**. **DO NOT** get brace wet. Brace can be removed for bathing and exercises **ONLY**.

WEIGHTBEARING STATUS

You will be **PARTIAL WEIGHTBEARING** on your operative leg. Use crutches or a walker **AT ALL TIMES**. Brace **NEEDS** to be on and locked in extension. **DO NOT** engage in prolonged periods of standing or walking over the first 7-10 days following surgery. Elevate the operative leg to chest level whenever possible to decrease swelling.

COLD THERAPY MACHINE

If provided, begin using your cold therapy machine after surgery. We recommend use at least 3 times per day, following the manufacturers instructions for setup and utilization of water and ice, for **NO LONGER THAN 45 MINUTES** per treatment. **ALWAYS** make sure to apply a barrier between the cooling pad and your skin. If you have not been provided with a cold therapy machine, apply ice to the surgical area in a waterproof bag for **NO LONGER THAN 30 MINUTES** per treatment.

DVT (BLOOD CLOT) PREVENTION

- >> Beginning the day of surgery, take one **ASPIRIN 81MG** tablet, twice per day for 4 weeks unless not allowed for other medical reasons. **Clear this with your family doctor before starting the aspirin.**
- >> Perform ankle range of motion exercises (ankle pumps) on both legs to exercise the calf muscles. Do ten reps every hour minimum while awake.
- >> **You should not do any flying for 2 weeks after surgery**

PAIN MEDICATION REFILL PROTOCOL

You **MUST** give your physician **48 HOURS** notice for any medication refill. Many medications require your surgeon to send the medication electronically and cannot be called into the pharmacy. Prescriptions **WILL NOT** be filled over the weekend. A prescription for the same medication you have been taking after surgery **IS NOT GUARANTEED**.

WHEN TO CALL YOUR SURGEON

Complications after surgery are fortunately very rare. Please contact our office if you have any of the following symptoms:

- >> Persistent fever >101°F, chills
- >> Increasing pain at the surgical site

- >> Increased swelling/pain in the calf
- >> New onset numbness or tingling
- >> Hives or new rashes
- >> Shortness of breath or chest pain
- >> Persistent nausea/vomiting
- >> Drainage from your incision lasting >5 days